arXiv:1305.0182v1 [stat.ME] 1 May 2013

Space-filling Latin Hypercube Designs based on Randomization Restrictions in Factorial Experiments

Pritam Ranjan* and Neil Spencer
Department of Mathematics and Statistics, Acadia University, Wolfville, Canada, B4P2R6
*Corresponding author: Pritam Ranjan, email: pritam.ranjan@acadiau.ca

Abstract

Latin hypercube designs (LHDs) with space-filling properties are widely used for emulating computer simulators. Over the last three decades, a wide spectrum of LHDs have been proposed with space-filling criteria like minimum correlation among factors, maximin interpoint distance, and orthogonality among the factors via orthogonal arrays (OAs). Projective geometric structures like spreads, covers and stars of $PG(p-1,q)$ can be used to characterize the randomization restriction of multistage factorial experiments. These geometric structures can also be used for constructing OAs and nearly OAs (NOAs). In this paper, we present a new class of space-filling LHDs based on NOAs derived from stars of $PG(p-1,2)$.

Keywords: Computer experiments; Nearly orthogonal arrays; Spreads; Stars.

1 Introduction

Latin hypercube sampling scheme was first proposed by McKay et al. (1979) as an alternative to random sampling in the Monte Carlo methods for numerically integrating complex multidimensional functions. Later on, the Latin hypercube designs (LHDs) became very popular in computer experiments for building statistical metamodels (Santner et al. 2003). Random LHDs can easily be constructed and are also useful in some applications, for instance, numerical integrations. However, random LHDs are not always suitable from a modeling viewpoint, for example, if all points are aligned along the main diagonal of the input space.

Space-filling designs have become very popular in computer experiments for building surrogates of the deterministic simulators. Since replicate runs of a deterministic simulator generate identical outputs, it is preferred that the inputs are spread out in the design space. Popular space-filling designs proposed thus far include, A, D, I-optimal designs (for minimizing generalized variance), mean squared error based designs, distance based designs (maximin and minimax), uniform designs, space-filling LHDs, orthogonal arrays (OAs), scrambled nets and Sobol sequences (see Santner et al. (2003) and Fang et al. (2006) for an overview).

In this paper, we discuss space-filling LHDs, a popular class of designs in computer experiments. Over the last three decades, a wide spectrum of LHDs have been proposed with different space-filling criteria, for instance, minimum correlation among factors (Iman and Conover 1982), maximin interpoint distance (Morris and Mitchell 1995), and orthogonality among the factors via OAs (Owen 1992; Tang 1993). The construction of LHDs with space-filling criteria like maximin distance or minimum correlation often requires computationally intensive search, whereas, OA-based LHDs are easy to construct as long as the OAs exist.

The existence of a desired OA is not always guaranteed, and the construction can also be challenging (Hedayat et al. 1999). OAs can be constructed using a variety of combinatorial objects like linear codes, difference schemes and mutually orthogonal Latin squares. Rains et al. (2000) discussed the existence and construction of OAs using a spread of a finite projective geometry, $\mathcal{P} = PG(p-1, q)$, and called them *geometric OAs*. A *spread* of $\mathcal{P}$ is a set of disjoint subspaces or flats of $\mathcal{P}$ that cover $\mathcal{P}$. Ranjan et al. (2009) established the equivalence between 2^p factorial experiments with multiple randomization restrictions and different geometric structures of $PG(p-1, 2)$ (e.g., spreads and covers). Here, a randomization defining contrast subspace (RDCSS) with $t(\leq p)$ independent randomization factors is equivalent to a $(t-1)$-flat of $PG(p-1, 2)$. This paper focuses on a class of LHDs that are based on arrays constructed using RDCSSs of a 2^p factorial experiment (i.e., we only consider $q = 2$).

For efficient analysis of a multistage factorial experiment, it is desirable to construct disjoint RDCSSs. Ranjan et al. (2009) established the existence of a desired set of disjoint RDCSSs via the existence of a spread of a $PG(p-1, 2)$. However, in many practical situations (e.g., the plutonium alloy experiment of Bingham et al. 2008), overlap among the RDCSSs cannot be avoided. For such cases, Ranjan et al. (2010) proposed designs based on a new geometric structure called a *star* - a set of distinct flats of $PG(p-1, q)$ that share a common overlap (the nucleus). That is, the proposed design is a set of distinct RDCSSs with a common overlap. A star that is also a cover (*covering star*) of $PG(p-1, q)$ simplifies to a spread if the nucleus is empty.

We have discovered a new class of space-filling LHDs that can be constructed using stars of $PG(p-1, 2)$. It turns out that a star with non empty nucleus generates nearly OAs (NOAs). In the spirit of Rains et al. (2000), we refer to these star-based NOAs as *geometric NOAs*. By following Tang's OA-based LHD construction algorithm, we construct a class of geometric NOA-based LHDs. Although such LHDs are not always very space-filling, a near orthogonality (e.g., Xu and Wu 2001) or space-filling criterion can be used to search for a good one. To avoid the search, we propose a set of guidelines for carefully distributing the factorial effects among RDCSSs of the star which ensures space-filling LHDs. Moreover, the existence of OA-based LHDs are limited to only few $n \times d$ combinations, whereas, the existence conditions for stars are less stringent.

The remainder of the paper is organized as follows. Section 2 presents an overview of LHDs, RDCSSs in a 2^p factorial experiment, and spreads and stars of a $PG(p-1, q)$. In Section 3, we establish theoretical results for the existence and an algorithm of the construction of geometric-NOAs. Section 4 concludes the paper with a few remarks.

2 Background

This section starts with a brief review of random LHDs and OA-based LHDs. Then a few results are presented to establish the equivalence between a multistage factorial design with randomization restrictions and geometric structures of $PG(p-1, 2)$.

2.1 Latin hypercube designs

Let $L(n, d)$ be an LHD with n runs in d factors, where L_{ij} denote the level of factor j in the i-th experimental run, and each factor includes n uniformly spaced levels. In computer experiments, the input spaces are typically bounded hyperrectangles, and can be transformed to unit hypercubes. A random $L(n, d)$ in $[0, 1]^d$ has $L_{ij} = (\pi_j(i) - u_{ij})/n$ for $1 \leq j \leq d$ and $1 \leq i \leq n$, where $u_{ij} \sim Unif(0, 1)$ and $\{\pi_j(1), ..., \pi_j(n)\}$ is a random permutation of $\{1, ..., n\}$ (see Tang (1993) for details). As a result, there are $(n!)^d$ distinct LHDs, even if we ignore the slight variation due to $Unif(0, 1)$ perturbations.

Although LHDs have nice one-dimensional projection property, that is, one point each in $((i-1)/n, i/n)$ for $1 \leq i \leq n$, random LHDs can be quite undesirable from modeling viewpoint. Figure 1 presents two realizations of random LHDs in $[0, 1]^2$.



(a) A good design for modeling (b) A bad design for modeling

Figure 1: Two realizations of random LHDs in $[0, 1]^2$.

Space-filling LHDs are commonly used in computer experiments for building statistical surrogates of complex computer simulators. In this paper, we propose star-based LHDs which are generalizations of the OA-based LHDs.

Definition 1 *An* $N \times d$ *array* A *denoted by* $OA(N, s_1 s_2 \cdots s_d, r)$ *is said to be a strength* r *OA with* N *runs and* d *factors, if factor* j *has* s_j *levels* $\{0, ..., s_j - 1\}$ *and each* $N \times r$ *subarray contains every possible* r*-tuple an equal number of times.*

The special case of $s_1 = s_2 = \cdots = s_d$ corresponds to a symmetric OA denoted by $OA(N, s, d, r)$. Although mixed-level (or asymmetric) OAs have been investigated in recent years (see Hedayat et al. (1999) for references), it is a relatively much less explored area as compared to the symmetric OAs.

A simple existence condition of an asymmetric OA follows from the strength aspect of Definition 1, that is, N must be a multiple of $s_1^{u_1} s_2^{u_2} \cdots s_p^{u_d}$ for every set of $u_1, ..., u_d \in \{0, 1\}$ such that $\sum_{i=1}^{d} u_i \leq r$. Another popular existence result comes from the Rao bound: $N - 1 \geq \sum_{i=1}^{d}(s_i - 1)$. Determining the existence of OAs is nontrivial, and becomes more and more difficult as the strength r and the number of levels s_j increase. Even OAs with strength 2 do not always exist for arbitrary N and d (see Hedayat et al. (1999) and Rains et al. (2002) for more results).

An $OA(N, s_1 s_2 \cdots s_d, r)$-based LHD will have N runs and d factors with $L_{ij} = (\pi_j(A_{ij}) + u_{ij})/s_j$ for $1 \leq j \leq p$ and $1 \leq i \leq N$, where $u_{ij} \sim Unif(0, 1)$ and π_j are independent random permutation of $\{0, ..., s_j - 1\}$. As a result, the input space $[0, 1]^d$ is partitioned in to a $s_1 \times s_2 \times \cdots \times s_d$ grid and N points are assigned to the cells according to the scheme directed by the rows of OA. Tang (1993) showed that LHDs based on a strength r OA inherit r-dimensional projection property. However, such LHDs do not guarantee projection property (i.e., space-filling) for lower dimensional hyperspace (see Figure 2).



Figure 2: An $OA(9, 4, 3, 2)$-based LHD in $[0, 1]^2$.

In this paper, we propose a new class of star-based geometric NOAs for constructing space-filling LHDs. As mentioned earlier, OAs exist for only a few $n \times d$ combinations, whereas, the existence conditions for stars are less stringent.

2.2 RDCSSs and Projective Geometries

Ranjan et al. (2009) proposed a unified framework using finite projective geometry for the existence and construction of multistage factorial designs with randomization restriction (e.g., nested designs, split-plot designs, split-lot designs, and combinations thereof). Let $\mathcal{P} = PG(p-1, 2)$ be the $(p-1)$-dimensional finite projective space defined by the set of all p-dimensional pencils over $GF(2)$. Here, a p-dimensional pencil with r nonzero elements uniquely corresponds to an r-factor interaction, and $\mathcal{P}$ denotes the set of all $2^p - 1$ factorial effects (excluding the grand mean).

The restrictions on the randomization of experimental runs lead to grouping experimental units into sets of trials. These sets are formed by using linearly independent pencils of $\mathcal{P}$, also referred to as the *randomization restriction factors* (like the blocking factors in a blocked factorial design). A set S of all non-null pencils formed as linear combinations of t independent randomization restriction factors in $\mathcal{P}$ constitutes a $(t-1)$-dimensional subspace (or, $(t-1)$-flat) of $\mathcal{P}$ (here $|S| = 2^t - 1$). We call such a subspace a t-dimensional randomization defining contrast subspace (RDCSS).

For efficient analysis of such experiments, it is desirable to construct disjoint RDCSSs. Ranjan et al. (2009) established the existence of a set of disjoint RDCSSs in a 2^p factorial experiment via the existence of a *spread* of a $PG(p-1, 2)$.

Definition 2 *For $1 \le t \le p$, a balanced $(t-1)$-spread of $\mathcal{P} = PG(p-1, q)$ is a set, ψ, of $(t-1)$-flats of $\mathcal{P}$ which partitions $\mathcal{P}$.*

The size of a balanced $(t-1)$-spread ψ of $PG(p-1, q)$ is $|\psi| = (q^p - 1)/(q^t - 1)$. A necessary and sufficient condition for the existence of a $(t-1)$-spread is that t divides p (André 1954). See Ranjan et al. (2009) for more results on the balanced and mixed (partial) spreads. In many practical situations (e.g., the plutonium alloy experiment of Bingham et al. (2008)), the overlap among the RDCSSs cannot be avoided. For such cases, Ranjan et al. (2010) proposed designs based on a new geometric structure called a *star* - a set of distinct flats of $PG(p-1, q)$ that share a common overlap (the nucleus).

Definition 3 *A balanced covering star $\Omega = St(\mu, t, t_0)$ of $\mathcal{P} = PG(p-1, q)$ is a set of μ rays ($(t-1)$- flats) and a nucleus ((t_0-1)- flat), where the nucleus is contained in each of the μ rays (i.e., $t_0 < t$), and $\mu = (q^{p-t_0} - 1)/(q^{t-t_0} - 1)$.*

A necessary and sufficient condition for the existence of a balanced covering stars $St(\mu, t, t_0)$ of $PG(p-1, q)$ is $(t-t_0)$ divides $(p-t_0)$. Let $St(t_1, ..., t_\mu; t_0)$ be a mixed or unbalanced covering star of $PG(p-1, q)$ with μ rays and (t_0-1)-dimensional nucleus, such that $t_1 \leq \cdots \leq t_\mu$. The next two lemmas are taken from Ranjan et al. (2010).

Lemma 1 *For the existence of a covering star* $St(t_1, ..., t_\mu; t_0)$ *of* $\mathcal{P} = PG(p-1, q)$, *the following conditions are necessary:*

(i) $q^{p-t_0} - 1 = \sum_{i=1}^{\mu}(q^{t_i-t_0} - 1)$,

(ii) $t_i + t_j - t_0 \leq p$ *for every* $i \neq j$ $(i, j = 1, ..., k)$,

A necessary and sufficient condition for unbalanced covering star is still unknown. The next result guarantees the existence of a balanced star for every t and p $(t < p)$.

Lemma 2 *For every* t $(2 \leq t < p)$ *and* $t_0 = t - 1$, *there exists a balanced covering star* $St(\mu, t, t_0)$ *of* $\mathcal{P} = PG(p-1, q)$ *with* $\mu = (q^{p-t+1} - 1)/(q - 1)$.

Next we discuss how these star-based multistage factorial designs can be used to construct space-filling LHDs.

3 Star based LHD

In this section, we first establish new existence results for geometric NOAs that are derived from stars. Then we propose an algorithm (a generalization of Rains et al. (2002)) for constructing such NOAs. Finally, we present a few guidelines for constructing NOAs that lead to space-filling LHDs.

Theorem 1 *The existence of a covering star* $St(t_1, ..., t_\mu; t_0)$ *of* $PG(p-1, 2)$, *is a sufficient condition for the existence of an* $NOA(2^p, s_{i_1}s_{i_2}\cdots s_{i_k}, 2)$ *with* $1 \leq k \leq \mu$, $1 \leq i_1 < i_2 < \cdots < i_k \leq \mu$ *and* $s_j = 2^{t_j}$ *for all* $1 \leq j \leq \mu$.

Similarly, the existence of a balanced star $St(\mu; t; t_0)$ of $PG(p-1, 2)$ suffices the existence of $NOA(2^p, 2^t, k, 2)$ for all $1 \leq k \leq \mu$. Moreover, it turns out that the maximal $NOA(2^p, 2^t, \mu, 2)$ is a 2^{-t_0} fraction of an $OA(2^{p+t_0}, 2^t, \mu, 2)$, and its existence requires $(t-t_0)|(p-t_0)$ which follows from the existence of a balanced star.

Theorem 2 *For every* $1 \leq t < p$, *there exist* $NOA(2^p, 2^t, k, 2)$ *for all* $1 \leq k \leq 2^{p-t+1}-1$, *and the maximal* $NOA(2^p, 2^t, 2^{p-t+1}-1, 2)$ *is a* $2^{-(t-1)}$ *fraction of an* $OA(2^{p+t-1}, 2^t, 2^{p-t+1}-1, 2)$.

Proofs of Theorems 1 and 2 follows from Lemma 1 and 2, and the construction of such NOAs using covering stars of $\mathcal{P} = PG(p-1, 2)$. Algorithm 1 is a generalization of the spread to OA construction recipe presented in Rains et al. (2002). Note that a covering star with empty nucleus (i.e., $t_0 = 0$) is a spread.

For this construction, we use the binary (vector or pencil) representation of all effects. Let $\{a_1, a_2, ..., a_{2^p-1}\}$ be the ordered set of all effects in $\mathcal{P}$, and $St(t_1, ..., t_\mu; t_0)$ be a covering star of $\mathcal{P}$ with μ rays $\{R_1, ..., R_\mu\}$, where $|R_j| = 2^{t_j}$ ($= s_j$, say). Algorithm 1 constructs an $NOA(2^p, s_1 s_2 \cdots s_\mu, 2)$ denoted by $A = [A_{*1} : A_{*2} : \cdots : A_{*\mu}]$.

Algorithm 1 Star to NOA construction

```
for $j = 1 \to \mu$ do
    For $j$-th stage of randomization restriction, find $t_j$ independent randomization restriction factors $\{\delta_1^{(j)}, ..., \delta_{t_j}^{(j)}\}$ such that $R_j = \langle \delta_1^{(j)}, ..., \delta_{t_j}^{(j)} \rangle$.
    for $i = 1 \to 2^p - 1$ do
        for $l = 1 \to t_j$ do
            Compute $b_{lj}^{(i)} = a_i \cdot \delta_l^{(j)}$    # inner product over $mod(2)$
        end for
        Define $A_{ij} = \sum_{l=1}^{t_j} b_{lj}^{(i)} 2^{t_j - l}$.
    end for
end for
```

For every $j \in \{1, 2, ..., \mu\}$, the i-th element of the NOA is $A_{ij} \in \{0, 1, ..., 2^{t_j} - 1\}$ for all $1 \leq i \leq 2^p - 1$. As a convention, we append a row of zeros at the beginning. We now present a few examples to illustrate the theoretical results and Algorithm 1.

Example 1 Let Ω be a covering star of $PG(3, 2)$ (or, in a 2^4 experiment) such that the rays (or RDCSSs) are of size seven each (i.e., $p = 4$ and $t = 3$). Here, $\{a_1, ..., a_{15}\} = \{D, C, CD, B, ..., ABCD\}$. Since there always exists a covering star for $t_0 = t - 1$ (with $\mu = 2^{4-2} - 1 = 3$), Theorem 2 ensures the existence of $NOA(16, 8, k, 2)$ for all $1 \leq k \leq 3$. Following the construction in Ranjan et al. (2010), let $\Omega = \{R_1, R_2, R_3\}$ with $R_1 = \langle A, B, ACD \rangle$, $R_2 = \langle C, D, ABC \rangle$ and $R_3 = \langle AC, BC, AD \rangle$. Note that the nucleus is $\langle AB, CD \rangle$.

The first column of the maximal NOA is $A_{*1} = (0, A_{11}, A_{21}, ..., A_{15,1})^T$, with, $A_{i1} = b_{11}^{(i)} 2^{3-1} + b_{21}^{(i)} 2^{3-2} + b_{31}^{(i)} 2^{3-3}$. As an example, for $i = 2$, $a_2 = C$ and $b_{11}^{(2)} = (1, 0, 0, 0)^T \cdot (0, 0, 1, 0) = 0$, $b_{21}^{(2)} = (0, 1, 0, 0)^T \cdot (0, 0, 1, 0) = 0$ and $b_{31}^{(2)} = (1, 0, 1, 1)^T \cdot (0, 0, 1, 0) = 1$. Therefore, $A_{21} = 0 \cdot 4 + 0 \cdot 2 + 1 \cdot 1 = 1$. The transpose of the final NOA is as follows:

$$\mathbf{A^T} = \begin{pmatrix} 0 & 1 & 1 & 0 & 2 & 3 & 3 & 2 & 5 & 4 & 4 & 5 & 7 & 6 & 6 & 7 \\ 0 & 2 & 5 & 7 & 1 & 3 & 4 & 6 & 1 & 3 & 4 & 6 & 0 & 2 & 5 & 7 \\ 0 & 1 & 6 & 7 & 2 & 3 & 4 & 5 & 5 & 4 & 3 & 2 & 7 & 6 & 1 & 0 \end{pmatrix}.$$

Though LHDs derived from star-based NOAs have nice geometric features, not all such LHDs are very space-filling. One can use a near orthogonality (Xu and Wu 2001) or space-filling criterion to choose a suitable NOA for constructing space-filling LHD, however, the search may still be computationally intensive. To avoid the search, we propose a few guidelines (not exhaustive) for carefully distributing the factorial effects among the RDCSSs of the star which ensures space-filling LHDs.

- (G1) $\delta_1^{(j)}$ (the first randomization factor in any ray) is not an element of the nucleus
- (G2) $\delta_l^{(j_1)} \neq \delta_l^{(j_2)}$ for all l
- (G3) $\delta_{l_1}^{(j_1)} + \delta_{l_2}^{(j_1)} \neq \delta_{l_1}^{(j_2)} + \delta_{l_2}^{(j_2)}$ for all $(l_1, l_2) \in \{1, 2, ..., t_{j1}\} \times \{1, 2, ..., t_{j2}\}]$

These guidelines suggest a careful selection of the randomization restriction factors for constructing the desirable geometric NOA. In the next few examples we use the star in Example 1 but choose $\delta_l^{(j)}$'s carefully to illustrate the importance of these guidelines.

Example 2 Consider the same star as in Example 1, that is, a covering star $\Omega = St(3;3;2)$ of $PG(3,2)$ (with $p = 4$, $t = 3$ and $t_0 = 2$), however, choose $\delta_l^{(j)}$'s such that only "G1" is violated. Let $R_1 = \langle AB, B, ACD \rangle$, $R_2 = \langle D, C, ABC \rangle$ and $R_3 = \langle AC, BC, CD \rangle$. Note $\delta_1^{(1)} = AB$ belongs to the nucleus. Figure 3 shows the corresponding LHDs with identity permutation and no $Unif(0,1)$ perturbation.



(a) A bad design (R_1 and R_2) (b) A bad design (R_1 and R_3) (c) A good design (R_2 and R_3)

Figure 3: Three realizations of $NOA(16, 8, 2, 2)$-based LHDs in $[0,1]^2$.

It is clear from Figure 3 that LHDs derived using R_1 (Figures 3(a) and 3(b)) are not as space-filling as compared to that obtained without it (Figure 3(c)).

Example 3 Consider the same star as in Examples 1 and 2, however, choose $\delta_l^{(j)}$'s such that only "G2" is violated. Let $R_1 = \langle A, B, ABCD\rangle$, $R_2 = \langle C, D, ABCD\rangle$ and $R_3 = \langle AC, BD, BC\rangle$. Note $\delta_3^{(1)} = \delta_3^{(2)} = ABCD$, which also belongs to the nucleus. Figure 4 shows the corresponding LHDs with identity permutation and no $Unif(0,1)$ perturbation.



(a) A bad design (R_1 and R_2) (b) A good design (R_1 and R_3) (c) A good design (R_2 and R_3)

Figure 4: Three realizations of $NOA(16, 8, 2, 2)$-based LHDs in $[0, 1]^2$.

It is clear from Figure 4(a) that LHD based on (R_1, R_2) shows correlation among the two factors (induced due to $\delta_3^{(1)} = \delta_3^{(2)} = ABCD$), and may not be very desirable for modeling. LHDs shown in Figures 4(b) and 4(c)) are reasonably space-filling in two-dimensional space.

Example 4 Consider the same star as in Examples 1 - 3, however, choose $\delta_l^{(j)}$'s such that only "G3" is violated. Let $R_1 = \langle A, B, ACD\rangle$, $R_2 = \langle C, ABD, ABC\rangle$ and $R_3 = \langle AC, AD, BC\rangle$. Note that (R_1, R_3) violate "G3" as $\delta_2^{(1)} + \delta_3^{(1)} = B + ACD = ABCD$ and $\delta_2^{(3)} + \delta_3^{(3)} = AD + BC = ABCD$, and (R_2, R_3) violate "G3" as $\delta_1^{(2)} + \delta_3^{(2)} = C + ABC = AB$ and $\delta_1^{(3)} + \delta_3^{(3)} = AC + BC = AB$. Figure 5 shows the corresponding LHDs.



(a) A good design (R_1 and R_2) (b) An okay design (R_1 and R_3) (c) A bad design (R_2 and R_3)

Figure 5: Three realizations of $NOA(16, 8, 2, 2)$-based LHDs in $[0, 1]^2$.

It is clear from Figure 5 that the violation of "G3" caused adverse effect on the randomness of the points and space-filling behaviour of the LHD.

Example 5 Consider the same star as in Examples 1 - 4, however, we wish to choose $\delta_l^{(j)}$'s that follow all three guidelines (i.e., no violations). Let $R_1 = \langle B, ACD, AB\rangle$, $R_2 = \langle D, C, ABC\rangle$ and $R_3 = \langle AC, BC, CD\rangle$. Note that the set $\{R_1, R_2, R_3\}$ here is very similar to that in Example 2, except the order of $\delta_l^{(1)}$ (randomization restriction factors for R_1). Figure 6 depicts the corresponding LHDs.



(a) A good design (R_1 and R_2) (b) A good design (R_1 and R_3) (c) A good design (R_2 and R_3)

Figure 6: Three realizations of $NOA(16, 8, 2, 2)$-based LHDs in $[0, 1]^2$.

It is clear from Figure 6 that the NOA derived from this star generates LHD with two-dimensional space-filling projections.

4 Concluding Remarks

This paper presents a new class of space-filling LHDs that are based on geometric NOAs derived from covering stars – designs for multistage factorial experiments with randomization restrictions. Though we assumed two-level factorial designs (i.e., $PG(p-1, 2)$) in Section 3, theoretical results and construction algorithms can easily be generalized for mixed/unbalanced covering or partial stars in $PG(p-1, q)$.

Although we have proposed a few guidelines for constructing space-filling NOA-based LHDs, it may not always be possible to satisfy all conditions (G1, G2 and G3) in a star construction. In such a case, one can use a near orthogonality (Xu and Wu 2001) or space-filling criterion to choose a suitable NOA for constructing space-filling LHD.

In Example 1, the LHDs are based a covering star $St(3; 3; 2)$ of $PG(3, 2)$, i.e., we chose $t = p - 1$. One could instead use another $2 \leq t < p$ for constructing a non-trivial star.

For instance, $t = 2$ and $t_0 = t - 1$ would generate a covering star $St(7; 2; 1)$ of $PG(3, 2)$. The advantage of using $St(7; 2; 1)$ instead of $St(3; 3; 2)$ would be the opportunity to construct LHDs with $d = 7$, however, the number of levels per factor will be reduced from eight to four. That is, the design would be slightly less space-filling in the one-dimensional hyperspace. Thus large covering stars are more useful.

Acknowledgments

We would like to thank Boxin Tang (Simon Fraser University) for many helpful comments and discussion. Ranjan's work was supported by Discovery grants from the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

André, J. (1954). Uber nicht-Desarguessche Ebenen mit transitiver Translationsgruppe. *Math. Z.*, 60, 156–186.

Bingham, D., Sitter, R., Kelly, E., Moore, L., and Olivas, J. D. (2008). Factorial designs with multiple levels of randomization. *Statist. Sinica*, 18, 493–513.

Fang, K.-T., Li, R. and Sudjianto, A. (2006). *Design and Modeling for Computer Experiments*. Chapman and Hall/CRC. Boca Raton, FL.

Hedayat, A. S., Sloane, N. J. A. and Stufken, J. (1999). *Orthogonal Arrays: Theory and Applications*. New York: Springer-Verlag.

Iman, R.L. and Conover, W.L. (1982). A distribution-free approach to inducing rank correlation among input variables. *Communication in Statistics - Simulation and Computation*, 11, 311–334.

McKay, M. D., Beckman, R. J. and Conover, W. J. (1979), A comparison of three methods for selecting values of input variables in the analysis of output from a computer code, *Technometrics*, 21(2), 239–245.

Morris, M.D. and Mitchell, T.J. (1995), Exploratory Designs for Computational Experiments, *Journal of statistical planning and inference*, 43, 381-402.

Owen, A.B. (1992). Orthogonal arrays for computer experiments, integration and visualization, *Statistical Sinica*, 2, 439-452.

Rains, E.M., Sloane, N.J.A., and Stufken, J. (2002). The lattice of N-run orthogonal arrays. *J. Statist. Plann. Inf.* 102, 477–500.

Ranjan, P., Bingham, D. and Dean, A. (2009), Existence and Construction of Randomization Defining Contrast Subspaces for Regular Factorial Designs, *The Annals of Statistics*, 37, 3580 – 3599.

Ranjan, P., Bingham, D. and Mukerjee, R. (2010), Stars and Regular Fractional Factorial Designs with Randomization Restrictions, *Statist. Sinica*, 20, 1637-53.

Santner, T.J., Williams, B. and Notz, W. (2003). *The Design and Analysis of Computer Experiments.* Springer Verlag, New York.

Tang, B. (1993). Orthogonal Array-Based Latin Hypercubes. *Journal of the American Statistical Association*, 88, 1392–1397.

Xu, H. and Wu, C. F. J. (2001), Generalized minimum aberration for asymmetrical fractional factorial designs, *The Annals of Statistics*, 29, 1066–1077.